



Draw the Triceratops






Triceratops horridus

Late Cretaceous, USA

Length: 9m;Weight: 11,000kg

As heavy as a truck, with an enormous neck firll and three impressive homs sprouting from its face, *Tricerotops* was a powerful beast. It had one of the largest skulls of any land animal and is thought to have lived in herba.







Dromaeosauridae

A family of fast-running, agile, feathered carnivores, dromaeosaurids first appeared in the Middle Jurassic but really flourished in the Cretaceous period, when they spread across the globe. They all shared a characteristic S-shaped neck and they had long arms that in some species could be folded against the body, as in the wings of modern birds.







Maiasaura peeblesorum

Late Cretaceous, USA Length: 9m; Weight: 3,000kg

This dinosaur's name means 'good mother lizard' because it was found caring for its eggs and young, Maissurar was a herbivore, feeding on plants, leaves, bernies and rotting wood. It roamed the Cretaceous plains in vast herds before returning to its nesting sites.





Coelophysis bauri



Sinosauropteryx prima

Ceratosaurus nasicornis



Diplodocus carnegii

Late Jurassic, USA
Length: 22–35m: Weight: 18,000kg
Diplodocus was one of the longest known dinosaurs,
in large part because of its whip-like tall, which it may
have used to defend itself. A Diplodocus could swing its
tall fast enough to create a loud cracking noise
to warn off predetors.





Troodon formosus

Late Cretaceous, North America Length: 2m; Weight: 50kg Troodon formosus had long hindimbs and large, retractable sickle-shaped claws on its second toes. It belonged to a family of dinosaurs thought to be the ancestor of living birds.







Stegosaurus armatus

Late Jurassic, USA and Portugal Length: 9m;Weight 2,300kg Stegosorurs comtus was the largest of the stegosaurs. It had diamond-shaped plates along its arched back and a long, narrow skall. It is thought to have eaten low-growing wegetation, including ferns, mosses and cycard plants.

How to draw a sauropod





Try it yourself



Riojasaurus incertus


Late Trassic Argentina
Late Trassic Argentina
The first true isauropods evolved at the end of the
Trassics and survived for more than 100 million years
until the end of the Cretacous. They include the largest
land animals ever to have lived and at their heaviest
land animals ever to have lived and at their heaviest
weighed close to 100,000(g. A key feature was their
long necks, which enabled them to outcompete other
herbivores by accessing a vide range of food sources.





Ankylosaurus magniventris

Late Cretaceous, USA and Canada Length: 7m; Weight: 3,00kg Ankylosourus magniventrs was the largest of the ankylosourus magniventrs was the largest of the ankylosourus, with horns on its head, a beak and small, leaf-shaped teeth. Some of its plates were fused together, giving its armour-plating extra strength: it would have needed it to see off the predators of the day, which included "promosurus rex."






Carcharodontosaurus saharicus skull

Mid-Cretacous, Africa
Length: 13m:Weight: 6,000kg
Carcharodortosours means jagged-toothed lizard and its
skull shows why—it was full of enormous senated teeth
Despite being one of the largest land predators of all time,
Carcharodortosourus would not have been without rivals.
It lived in the same region as the 18m-long \$pinosourus,
one of the mightest beats of the Mesozoic erus,









Tenontosaurus tilletti (being attacked by a pack of Deinonychus)

Early Cretaceous, North America Length: 8m; Weight: 1,500kg A close relative of *Iguanodon*, this ornithopod would have been capable of walking both on all fours or on its hind legs. Many Tenontosaurus fossils have been found with those of the bird-like theropod, Deinonychus, and a juvenile Tenontosaurus has even been found with Deinonychus bite marks in its bones.







Brachiosaurus altithorax

Late Jurasic, North America Length: 25m:Weight 28,000Kg
When its fossils were discovered in 1900, Berchissarurs took the record of the largest dinosure. Although it has now been surpassed, it is still one of the tallest known dinosaurs. It had a graffel-like body with a long neck and unusually long front limbs. Brachissarurs would have used its long neck to reach leaves, as it would have been unable to rear up on its infollimbs.

Which silhouette matches this Spinosaurus?




\mathcal{A} nswers



Spinosaurus aegyptiacus

Mid-Cretaceous, Egypt
Length: 18m; Weight 9,000kg
Spinosourus was one of the largest land predators ever.
However, unlike other big predators of the Cretaceous
period, its skull was long and narrow and full of conical
teeth that were ideal for spearing fish.






Sordes pilosus

Late Jurassic, Kazakhstan Wingspan: 0.6m; Weight: 5kg


Perosaurs survived from the Late Trassic to the end of the cretaceous pend and their fossis have been found on every continent. They evolved into a diverse range of species, from the size of a pigeon to that of a small plane. Perosaur wings were made up of a memirane of skin, muscle and tissue, which stretched from the ankle bones to an extremely elongated fourth finger.





Saurolophus angustirostris

Late Cretaceous, Mongolia
Length: 12m;Weight: 3,300kg
Saurolophus angustrostris was the most common Asian hadrosaur.
For millions of years this group of duck-billed dinosaurs were
the world's dominant herbivores. They evolved from the
iguanodontids and spread throughout Late Cretaceous
Asia, America and Europe.







lchthyosaurus communis




Kronosaurus queenslandicus





1 - Mei long – length: 40cm
 2 - Troodon formosus – length: 2m
 3 - Ornithomirmus edmontonicus – length: 3.8m
 4 - Kentroscurus aethiopicus – length: 5m
 5 - Euoplocepholus trutus – length: 6m
 6 - Tyrannosaurus rex – length: 12m
 7 - Potzopottan moyorum – length: 37m





1. Triassic 252 – 201 million years ago



2. Jurassic 201 – 145 million years ago



3. Cretaceous 145 – 66 million years ago

3 145 - 66 million years ago







Ceratosaurus nasicornis

Late Jurassic, USA
Length: 7m: Weight: 700kg
Cerotosourus was a medium-zized therepool distinguished
by a large horn over its note and two horn-like ridges
above its eyes. Some scientists think the horn was used by
males to fight over females, while others argue it was
used for display purposes only.



Dilong paradoxus

Early Cretaceous, China Length: Zm; Weight: 10kg Dillong was the first tyranoscaur found with evidence of primitive hair-like structures, which resembled flutfy, insulating down. Its discovery proved that it wasn't just the small, bird-like dimosaurs that possessed feathers.



Pentaceratops sternbergii

Late Cretaceous, USA
Length: 6.5m; Weight: 5,000kg
Pentocerotops stembergis name means: five horned
face' after the homs on its snout, eyebrows and cheeks.
Its skull was up to 3m long, making it the longest skull
of any land animal in history.

BONNIER

















Ferns and leaves



Ceratosaurus nasicornis




Kentrosaurus aethiopicus and other ornithopods




Spinosaurus aegyptiacus



Fish and other marine life



Copy these oviraptor heads





Gigantoraptor erlianensis


Late Cretaceous, Mongolia Learnier, H. (ADOkg This massive oviraptor stood as full as a graffe and was 35 times bigger than Citipoti, the next largest in the genus. It is the largest known beaked dinosaur and if it was feathered it would be one of the largest known feathered animas of all time.



Anzu wyliei

Late Cretaceous, North America
Length: 3m; Weight: 225kg
Nicknamed the chicken from hell Anzu wylele is the most
complete oviraptorosaur to be found outside Asia. Small prongs
of bone found on the roof of its mouth match those
found in today's egg-eating makes.

How to draw a plesiosaur




Try it yourself



Elasmosaurus platyurus

Late Cretaceous, USA Length: 14m; Weight: 2,000kg In the Mesozoic era, the seas and oceans were filled with a huge variety of large marine reptiles, including the plesiosaurs. They had two pairs of flippers and had extremely long necks, small heads and sharp, pointed teeth for catching fish.



Create your own duck-billed dinosaur





Tsintaosaurus spinorhinus



Parasaurolophus walkeri



Lambeosaurus lambei






Tyrannosaurus rex







 ${\color{blue}\textbf{Colour the pachycephalosaur heads}}$





Stygimoloch spinifer





Dracorex hogwartsia




Pachycephalosaurus wyomingensis


Match these species to the regions where their fossils have been found








- 1. Dilong paradoxus
- 2. Spinosaurus aegyptiacus
- 3. Kronosaurus queenslandicus
- 4. Kentrosaurus queensianaicu
- 5. Iguanodon bernissartensis
- Iguanoaon bernissariensis
 Patagotitan mayorum
- 7. Triceratops horridus





Spinosaurus aegyptiacus

Mid-Cretaceous, Egypt
Length: Bim-Weight 90,000kg
Instantly recognisable by the huge sail on its back, Spinosourus'
strange and specialized form meant that it was able to carve
out a niche for itself alonguide the other theropods of the day,
hunting for fish and other aquatic and shore-dwelling prey,
Many now believe its huge sail was used for display,
to wan off predators and to attract mates.







lchthyosaurus communis



 $Volaticotherium\ antiquum$

Colour in this Tyrannosaurus rex





Tyrannosaurus rex

Late Cretaceous, North America
Length: 12m;Weight: 6,000kg
The most famous dinosaur, first discovered in 1902,
Tyranosaurus has captured the imagination like no other.
With 60 teeth, a powerful bite and a superb sense of smell.
Tyranosaurus was undoubtedly a ferocious siller: Their short,
deep skulls allowed them to generate extreme bite forces,
which more than made up for their tiny but strong forelimbs.


Which silhouette matches this *Cynognathus?*















${\mathcal A}$ nswers














Cynognathus crateronotus

Early-Mid-Triassic, Africa, South
America and Antarctica
Length: 12m Weight 6.5kg
A fast, fierce predator, this small cynodont was heavily-built
with wide jaws and sharp teeth capable of slicing through
flesh. It was possibly warm-blooded and covered in hair:





Quetzalcoatlus northropi


Late Cretaceous, USA
Wingspan: 12m; Weight: uncertain
Quetzaicodulus was one of the largest animals ever to fly.
It had a huge toothless beak and a wingspan the length of a bus.
The strength in its thickened arm bones enabled it to fly and it is
thought to have taken off from a crouching position on all flours.




Dimorphodon macronyx

Early Jurassic, England

Dimorphodon's large clawed hands enabled this pterosaur
to climb up steep cliffs or it could then launch itself off and fly
It had both fangs and grinding teeth, suggesting it lived off
a diet of insects and small animals rather than fish.



Draw the *Plateosaurus*





Plateosaurus engelhardti

Late Trisser, Cermany, Switzerland and France Length: 10m; Weight: 4,000kg Ploteosourus engelhords was one of the best known European dinosaurs: Hundreds of their fossis have been found together in one place, suggesting that they lived in herds. They would have browsed for food among the high tree branches, rearing up on their back legs to reach the best foliage.



